

# FUNDAMENTALS OF PLANT PATHOLOGY

## PRACTICAL MANUAL

Course No. HPP-226

*For*

B.Sc. (Hons.) Horticulture 3(2+1)



Dr. Shubha Trivedi  
Dr. P. P. Jambhulkar  
Dr. Meenakshi Arya

2020

College of Horticulture & Forestry  
Rani Lakshmi Bai Central Agricultural University  
Jhansi-284003

**Syllabus HPP 226 3(2+1):** Familiarity with general plant pathological laboratory and field equipments. Study of disease symptoms and signs and host parasite relationship. Identification and isolation of plant pathogens. Koch's postulates. Preparation of fungicidal solutions, slurries, pastes and their applications.

**Name of Student** .....

**Roll No.** .....

**Batch** .....

**Session** .....

**Semester** .....

**Course Name :** .....

**Course No. :** .....

**Credit** .....

**Published: 2020**

**No. of copies: .....**

**Price: Rs.**

---

### CERTIFICATE

This is to certify that Shri./Km. ....ID No.....has completed the practical of course.....course No. .... as per the syllabus of B.Sc. (Hons.) Agriculture/ Horticulture/ Forestry ..... semester in the year.....in the respective lab/field of College.

Date:

Course Teacher

## INDEX

<b>S. No.</b>	<b>Title</b>	<b>Page No.</b>
1.	To get familiar with general plant pathological laboratory equipments	
2.	Parts of the Microscope, its parts and handling	
3.	Collection and preservation of plant disease samples	
4.	Preparation of Potato Dextrose Agar Medium	
5.	Isolation and purification of plant pathogens from diseased Plant Tissues	
6.	Demonstration of Koch's postulates	
7.	Identification of different types of mycelium and other fungal structures	
8.	Identification of different types of asexual fruiting bodies, sexual spores and ascocarps.	
9.	Identification of plant disease symptoms	
10.	Identification of the plant pathogens of Phylum Chytridiomycota and Oomycota	
11.	Identification of the plant pathogens of Phylum Zygomycota	
12.	Identification of the plant pathogens of Phylum Basidiomycota	
13.	Identification of the plant pathogens of Phylum Ascomycota	
14.	Staining and identification of plant pathogenic bacteria	
15.	To demonstrate Sap transmission of plant viruses	
16.	To identify Phanerogamic plant parasites.	
17.	To get familiar with different fungicides and their formulations	
18.	Calculation of fungicide sprays concentrations	
19.	Methods of application of fungicides and their safe use	

## Practical No. 1

### Objective: To get familiar with general plant pathological laboratory equipment

The students in batches will visit the laboratory of Plant Pathology to acquaint with different appliances, tools, glass-wares, and other miscellaneous items, which they will be using in various exercises and experiments to be conducted.

#### 1. Identify the laboratory equipments available in the Plant Pathology Laboratory:

##### (a) Laboratory appliances / tools:

(i)		(ii)	
(iii)		(iv)	
(v)		(vi)	
(vii)		(viii)	
(ix)		(x)	
(xi)		(xii)	
(xiii)		(xiv)	
(xv)		(xvi)	
(xvii)		(xviii)	
(xix)		(xx)	

##### (b) Glass-wares:

(i)		(ii)	
(iii)		(iv)	
(v)		(vi)	
(vii)		(viii)	
(ix)		(x)	
(xi)		(xii)	

**2. Label the following laboratory equipments and state its principle and functions.**

Auto Clave: .....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



Laminar Air Flow: .....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



BOD Incubator: .....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



Hot Air Oven: .....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



## Practical No. 2

**Objective: To get familiar with Microscope, its parts and handling**

**1. Draw a well labeled diagram of a Compound Microscope and indicate all the important parts.**


















**Practical No. 8**

**Objective:** Identification of different types of asexual fruiting bodies, sexual spores and ascocarps.

Identify different asexual fruiting bodies and ascocarps provided in the slides and draw the structures observed under the microscope and describe its characteristics.

Ascocarps:.....  
.....  
.....  
.....  
.....














.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....





.....



2. Record the characteristic differences in morphology of *Pythium* and *Phytophthora* and draw a neat and labeled diagram of the spores.

Characteristics	<i>Pythium</i> spp	<i>Phytophthora</i> spp
Mycelium	.....	.....
	.....	.....
	.....	.....
Sporangiophores	.....	.....
	.....	.....
	.....	.....
Sporangia	.....	.....
	.....	.....
	.....	.....
Oospores	.....	.....
	.....	.....
	.....	.....
Haustoria	.....	.....
	.....	.....
	.....	.....
Vesicle	.....	.....
	.....	.....
	.....	.....
Zoospore formation	.....	.....
	.....	.....
	.....	.....



3. State the systematic position of the Genera given in the space below. Record the characteristic morphology of Genus – *Peronospora* (Downy mildew), *Sclerospora* and draw a neat and labeled diagram of the spores along with conidiophores.

SYSTEMATIC POSITION	SYSTEMATIC POSITION
.....	.....
.....	.....
.....	.....
.....	.....
.....	.....
.....	.....

Characteristics	<i>Sclerospora</i>	<i>Peronospora</i>
Mycelium	.....	.....
	.....	.....
	.....	.....
Conidia	.....	.....
	.....	.....
	.....	.....
Branching	.....	.....
	.....	.....
Sterigmata	.....	.....
	.....	.....
Oospores	.....	.....
	.....	.....
	.....	.....
Conidiophores	.....	.....
	.....	.....
	.....	.....

Diagram	Diagram

4. Record the characteristic morphology of *Albugo candida* (White blister/rust) and draw a neat and labeled diagram of spores.

**SYSTEMATIC POSITION:**

.....

.....

.....

.....

.....

.....

<b>Characteristic</b>	<b>Description</b>
Mycelium	..... ..... .....
Sporangiophores	..... .....
Sporangia	..... .....
Oospores	..... ..... .....

Diagram

**Objective: Identification of the plant pathogens of Phylum Zygomycota**

1. Record the characteristic morphology of Genus – *Mucor* (Bread mould) and *Rhizopus* and draw a neat and labeled diagram of their spores.

**SYSTEMATIC POSITION**

.....

.....

.....

.....

.....

.....

**SYSTEMATIC POSITION**

.....

.....

.....

.....

.....

.....

Characteristics	<i>Mucor</i>	<i>Rhizopus</i>
Mycelium	.....	.....
	.....	.....
	.....	.....
Sporangiophores	.....	.....
	.....	.....
	.....	.....
Sporangia	.....	.....
	.....	.....
	.....	.....
Columella	.....	.....
	.....	.....
	.....	.....
Aplanospores	.....	.....
	.....	.....
	.....	.....
Zygosporangia	.....	.....
	.....	.....
	.....	.....

Diagram

Diagram


**Objective: Identification of the plant pathogens of Phylum *Basidiomycota***

Record characteristic morphology of the following Genera and draw a neat and labeled diagram of spores.

Genus: *Uromyces*

Features:

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....




.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

Genus: *Melampsora*

Features:

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....




.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

Genus: *Ustilago*

Features:

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....





**Objective: Identification of the plant pathogens of Phylum Ascomycota**

Class: Eurotiomycetes

<b>SYSTEMATIC POSITION (<i>Aspergillus</i>)</b> ..... ..... ..... ..... ..... .....	<b>SYSTEMATIC POSITION (<i>Penicillium</i>)</b> ..... ..... ..... ..... ..... .....
-------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------

<b>Characteristics</b>	<b><i>Aspergillus</i></b>	<b><i>Penicillium</i></b>
Mycelium	..... ..... .....	..... ..... .....
Foot Cell	..... ..... .....	..... ..... .....
Conidiophore	..... ..... .....	..... ..... .....
Vesicle	..... ..... .....	..... ..... .....
Sterigmata	..... ..... .....	..... ..... .....
Conidia	..... ..... .....	..... ..... .....
Perfect Stage	..... .....	..... .....

Diagram

Diagram

**Class: Sordariomycetes**

**SYSTEMATIC POSITION (*Fusarium*)**  
.....  
.....  
.....  
.....  
.....  
.....

**SYSTEMATIC POSITION (*Claviceps*)**  
.....  
.....  
.....  
.....  
.....  
.....

**Characteristics**

***Fusarium***

***Claviceps***

Mycelium	..... ..... .....	..... ..... .....
Sporodochia	..... ..... .....	
Conidiophore	..... ..... .....	
Conidia	..... ..... .....	..... ..... .....
Chlamyospores	..... ..... .....	
Sclerotia		..... ..... .....
Perfect Stage	.....	

Diagram

Diagram

**SYSTEMATIC POSITION (*Pyricularia*)**

.....

.....

.....

.....

.....

.....

**SYSTEMATIC POSITION (*Colletotrichum*)**

.....

.....

.....

.....

.....

.....

**Characteristics**

***Pyricularia***

***Colletotrichum***

Mycelium

.....

.....

.....

Conidiophore

.....

.....

.....

Conidia

.....

.....

.....

Acervuli

.....

.....

.....

Perfect Stage

.....

Dothideomycetes

**SYSTEMATIC POSITION (*Helminthosporium*)**

.....

.....

.....

.....

.....

.....

**SYSTEMATIC POSITION (*Alternaria*)**

.....

.....

.....

.....

.....

.....

**Characteristics**

***Helminthosporium***

***Alternaria***

Conidiophore

.....

.....

.....

.....

.....

.....

Conidia

.....

.....

.....

.....

.....

.....

Perfect Stage

.....

.....

--	--

**SYSTEMATIC POSITION (*Phyllosticta*)**

.....

.....

.....

.....

.....

.....

**SYSTEMATIC POSITION (*Cercospora*)**

.....

.....

.....

.....

.....

.....

**Characteristics**

***Phyllosticta***

***Cercospora***

Mycelium .....

.....

.....

Conidiophores .....

.....

.....

Pycnidia .....

.....

.....

Conidia .....

.....

.....

Perfect Stage .....

.....



**Class: Letiomycetes**

**SYSTEMATIC POSITION (*Erysiphe*)**

.....

.....

.....

.....

.....

.....

**SYSTEMATIC POSITION (*Sclerotinia*)**

.....

.....

.....

.....

.....

.....

**Characteristics**

***Erysiphe***

***Sclerotinia***

Mycelium

.....

.....

.....

Asexual Stage

.....

Conidiophores

.....

.....

.....

Conidia

.....

.....

.....

Sexual Stage

.....

Cleistothecia

.....

.....

.....

Apothecia

.....

.....

.....

Appendages

.....

.....

.....

Sclerotia

.....

.....

.....

Asci

.....

.....


.....

Ascospores

.....

.....

.....



**Class: Taphrinomycetes**

**SYSTEMATIC POSITION (*Taphrina*)**  
.....  
.....  
.....  
.....  
.....  
.....


**Characteristics**

***Taphrina***

Mycelium .....  
.....  
.....

Asci .....  
.....  
.....

Ascospores .....  
.....  
.....



**Objective: Staining and identification of plant pathogenic bacteria**

1. Prepare smear of given bacterial samples and perform gram-staining and identify on the basis of gram staining. Write the step by step procedures of gram-staining.

**Materials Required:** .....

.....

.....

**Procedure:**

**A. Smear preparation:** .....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**B. Gram-staining:** .....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Observation:**

Sl.No.	Color of the stain	Gram-reaction

**Objective: To demonstrate sap transmission of viruses**

Perform transmission of virus through sap using *tomato leaf curl virus* and note the symptoms.

**Materials Required:** .....

**Procedure:** .....

**Observations:** .....



**Objectives: To get familiar with different fungicides and their formulations**

1. Write the constituents of the following fungicides:

A. Bordeaux mixture: .....

.....  
.....  
.....  
.....

B. Bordeaux paste: .....

.....  
.....  
.....  
.....

C. Burgundy mixture: .....

.....  
.....  
.....  
.....

D. Cheshunt compound: .....

.....  
.....  
.....  
.....

E. Chaubattia Paste: .....

.....  
.....  
.....  
.....

Precautionary measures: .....

.....  
.....  
.....  
.....









**GENERAL PLANT PATHOLOGICAL LABORATORY EQUIPMENTS**

**(a) Laboratory appliances / tools:**

1. Autoclave	6. Hot-air oven	11. Scissor	16. Sprit Lamp
2. Freeze	7. Incubator	12. Cork-borer	17. Forceps
3. Hot Plate	8. Pan (different sizes)	13. Needle, Inoculating needle	18. Rotary shaker
4. Knife / Blade	9. Scalpel	14. Bearing Blander	19. Glass marker
5. Inoculating needles	10. Laminar flow	15. Gel electrophoresis	20. Centrifuge


**(b) Glass-wares:**

1. Conical flask (different sizes)	5. Beaker (different sizes)	9. Slides
2. Measuring cylinder (different capacity)	6. Pippet (different volume)	10. Watch glass
3. Petridishes	7. Culture tubes	11. Dropping bottle
4. Cover-slip	8. Nematode counting dish	12. Bearman funnel

**(c) Miscellaneous items:**

1. Cotton	5. Blotting paper	9. Washing brush
2. Aluminium foil	6. Wash bottle	10. Washing powder
3. Trays	7. Thread	11. Wire basket
4. Sieve of different sizes	8. Rubber bands	12. Mortar and pestle

**MICROSCOPE**



**Compound Microscope**

## Types of Microscopes:

1. Simple Microscope (Magnifying glass)
2. Student Compound Microscope
3. Stereoscopic Microscope
4. Electron Microscope
5. Simple dissecting Microscope Monocular
6. Compound Microscope (Binocular)
7. Phase Contrast Microscope

**Compound Microscope:** A Compound microscope consists of more than one lenses fitted one above the other at a proper distance (160mm) in a cylindrical tube. An object magnified by one lens (Objective) is further magnified by another lens (eye piece).

### Different Parts of a Compound Microscope

1. **Eyepiece:** It is a lens that fits into the top of body tube (drawtube). It is also called ocular lens. It is usually marked with 6X, 10X or 15X that mean it can magnify the object 6, 10 or 15 times.
2. **Drawtube:** This is a small cylindrical tube on the top of which eyepiece is fitted.
3. **Body tube:** It is a hollow cylindrical tube attached to upper end of the arm on which it can be moved up and down with the help of coarse adjustment knob.
4. **Arm:** It is a curved structure used for holding the microscope.
5. **Coarse adjustment knob:** This is used to locate the object by the objective.
6. **Fine adjustment knob:** Mostly fitted below coarse adjustment knob. It is used when the object is viewed either under high power or under low power for getting sharp and distinct view.
7. **Inclination Joint:** It is the point where microscope with stage and body with two lenses can be bent to a comfortable angle for smooth and strain free observation. This point lies close to the junction of stage and arm.
8. **Nose piece:** It is a disc like body fitted at lower end of the body tube. It has provisions for three lenses. It can be revolved also.
9. **Objectives:** They are the lenses of different magnifications, screwed in the nosepiece. Objective is also marked with 6X, 10X, 40X, 100X etc.
10. **Stage:** A flat rectangular or square plate with round aperture in the centre of the stage.
11. **Clip:** Two clips on either side of the aperture on the stage for holding the slide.
12. **Mechanical device:** Slide is fitted in it, which can be moved forward, backward, right and left to locate the object.
13. **Diaphragm:** It is a circular plate with several holes of different diameters and is attached underneath the stage. It can also be rotated so as to bring the hole of the diaphragm in front of the hole in the stage. It regulates the quantity of light towards body tube.
14. **Condenser:** It is used to regulate the intensity of light.
15. **Pillars:** These are the two vertical structures to give the support on the base of microscope.
16. **Mirror:** It is spherical reflecting mirror, which can be adjusted to direct the light through diaphragm, stage and lenses. It is fitted in mirror holder.
17. **Foot or Base:** It forms the base of the microscope.

**Precautionary measures:** 1. Closing one eye while using the other to look through a monocular microscope tends to tire certain eye muscles. Learn to keep both eyes open. 2. Another source of eyestrain results from imperfect focusing. Keep the hand on the fine adjustment and continually making slight changes in focus to study different parts of the field can obviate this. 3. Clean the lenses with tissue paper. 4. Never use ordinary tissue paper or cloth, which might contain grit. 5. Never rub the lenses heavily it may scratch the lens. 6. Do not remove lenses from their mounts or unscrew the objectives from the nosepiece.

## COLLECTION AND PRESERVATION OF PLANT DISEASE SAMPLES

Preservation means killing or restricting the growth of an organism in or on the substrate on which it grows.

### 1) Dry Preservation:

- a) **Collection and drying:** The sample should have distinctively visible symptoms. Dry the specimen in layer of blotting sheets under sunlight or in hot air oven for few days.
- b) **Labelling and packaging:** The material should be kept in good herbarium packets. This is attached to a chart paper sheets. The two sides of packet are folded first, then bottom flap and finally top flap. The name of pathogen, host, locality, date, name of scientist who identified the specimen, should be mentioned on the label.
- c) **Disinfection and storage:** The specimen folders are fumigated with methyl bromide vapours in fumigation chamber for 24-48 hrs before storage.

**2) Wet Preservation:** Washed fresh diseased specimens are put in a boiling mixture of 1 part of glacial acetic acid saturated with normal copper acetate crystals and 4 parts of water till the green colour reappears and then kept preserved in 5 per cent formalin in the glass jars. All mounted or preserved specimens must be labeled with as much of the following information as far as possible:

1. Host (name of the diseased plant)
2. Name of the disease Parasite (the name of the organism causing the disease)
3. Place where collected
4. Date of collection
5. Name of the collector

**Size of the specimen:** A specimen should ideally be 25–40 cm long and up to 26 cm wide, allowing it to fit on a standard herbarium mounting sheet which measures 42 x 27 cm. This is also the approximate size of tabloid newspapers. Plant parts that are too large for a single sheet may be cut into sections pressed on a series of sheets, for example a palm or cycad frond. Long and narrow specimens such as grasses and sedges can be folded once, twice or even three times at the time of pressing. In this way a plant of up to 1.6 m high may be pressed onto a single sheet. For very small plants, a number of individuals may be placed on each sheet.

### POTATO DEXTROSE AGAR MEDIUM


**Materials required:** Peeled potato slices (200g); Dextrose (20 g); Agar- agar (20 g); Distilled water (1000 ml)

**Method:**

- Potato slices are cooked in 500 ml of water.
- Then filtered with the help of muslin cloth.
- Agar-agar is melted in 500 ml of water.
- Potato juice is added to the melted agar.
- Volume is made 1000 ml by adding required water.
- Again, it is filtered through muslin cloth.
- Dextrose is added in this mixture and shaken well.
- Medium is sterilized in an autoclave at 1.1kg/cm<sup>2</sup> pressure for 20 minutes at temperature of 121.6°C. Thus the medium is ready for use.

### ISOLATION OF PLANT PATHOGENS FROM DISEASED PLANT TISSUES

Tissues sampled during the active stage of an infection are likely to have within them only the pathogen responsible for the infection; the surfaces of such tissues, however, are usually contaminated with saprophytic organisms. The steps of isolation of the pathogen have been given in the flowchart:



### KOCH'S POSTULATES

**Four steps of Koch Postulates:**


1. The suspected causal agent must be present in every diseased organism examined.
2. The suspected causal agent must be isolated from the diseased host organism and grown in pure culture.
3. When a pure culture of the suspected causal agent is inoculated into a healthy susceptible host, the host must reproduce the specific disease.
4. The same causal organism must be recovered again from the experimentally inoculated and infected host *i.e.*, the recovered agent must have the same characteristics as the organism in step 2. (The 4<sup>th</sup> postulates was appended by E.F.Smith)

### DIFFERENT STRUCTURES OF FUNGI

**Mycelium:** Network of hyphae is called as mycelium. It may be aseptate or septate.


**Aseptate Mycelium-** When the hyphae are undivided by cross-walls (septa) it is known as septate mycelium. This type of mycelium is found in lower fungi.

**Septate Mycelium-** When the mycelium is divided by cross walls (septa) at certain intervals, it is known as septate mycelium. In the septa (singular septum), there is a minute hole, which is known as "septal pore." This type of mycelium is found in higher fungi.



**Types of Asexual Spores:** Asexual spores are those in which sex is not involved. Generally, five types of asexual spores are produced in fungi.


1. **Arthrospores (Oidia):** Formed in chains (basipetal) on short conidiophores, single celled, barrel or drum shaped.
2. **Chlamydospores:** Formed singly or in chains, which may be terminal or intercalary, provided with an envelop (covering).
3. **Blastospores:** Spores formed by process of budding, which are single celled, first formed in chains but later separated from each other.
4. **Conidia:** Formed at the tip or side of the hypha (Conidiophore), may be formed singly or in chains, quite variable in shape, size, septation, colour and also in ornamentation.
5. **Zoospores:** Pear or kidney shaped, single shaped, naked, motile (flagellate), produced in sporangium (zoosporangium).
6. **Aplanospores:** Oval or spherical in shape, single celled, non-motile (aflagellate) and produced mostly in collumellate sporangium.



### TYPES OF ASEQUAL FRUITING BODIES, SEXUAL SPORES AND ASCOCARPS

#### Asexual fruiting bodies:


1. **Pycnidia:** These are spherical or flask shaped structures in which the conidia are produced. They have the natural opening known as ostiole through which the conidia are liberated. This type of structure is produced in order Spaeopsidales of sub division Deuteromycotina.
2. **Acervuli:** These are mat or cushion shaped structure formed below the cuticle or epidermis of the host. They may be provided with sterile hair like structures known as setae.
3. **Sporodochia:** These are the cushion-shaped structure on which the conidiophores are produced.
4. **Synnemata:** In these structures the conidiophores are grouped together at the base and free towards apex.



#### TYPES OF SEXUAL SPORES


Four types of sexual spores are formed in fungi, which are produced by various methods and they form the bases for the classification of fungi in different sub-divisions.

1. **Oospores:** Mostly spherical in shape, formed in the oogonium, usually smooth walled. They are formed by gametangial contact (oogamy), characteristics of phylum oomycota.
2. **Zygospores:** Black in colour, rough-walled, warty in appearance and provided with suspensors. They are formed by gametangial copulation (zygogamy), characteristics of sub-division Zygomycotina.
3. **Ascospores:** Produced in asci, definite in number (usually 8). They are formed by spermatization/ somatogamy, characteristics of sub-division Ascomycotina.
4. **Basidiospores:** Borne on the basidium, definite in number (usually 4). They are formed by spermatization/ somatogamy, characteristics of sub-division Basidiomycotina.



## TYPES OF ASCOCARPS

1. **Cleistothecia (-um):** Spherical in shape, black in colour, hard in structure and without any natural opening. Asci come out by tearing or breaking of the cleistothecium. Cleistothecia are also provided with appendages.
2. **Perithecia (-um):** Flask shaped with natural opening known as “**ostiole**”, sometime having long neck. Asci are produced in the perithecium at basal region. Paraphyses may also be present in between the asci.
3. **Apothecia (-um):** The ascocarp, which produces its asci in an open disc or cup shaped structure, is called as apothecium. It is exposed and form the layer of asci in a “**hymenium**” among them paraphyses may also be present.




## TYPES OF DISEASE SYMPTOMS PRODUCED DUE TO INFECTION BY PATHOGEN

1. **Blights:** A disease characterized by general and rapid killing of leaves, flowers and stems.
2. **Chlorosis:** When repression of colour is partial i.e., normally green tissues are yellow or when yellow colour is uniform and unbroken in leaves infected by plant pathogen.
3. **Mosaic:** Patches of normal green tissues alternate with yellow areas resulting in mottling, spotting, flecking, striping or blotching against the normal background tending to have a clearly defined boundary delineated by the veins.
4. **Vein-clearing:** is a kind of sub-type of mosaic where tissues close to veins turn yellow and remaining lamina surface remains green.
5. **Vein-banding:** is a kind of sub-type of mosaic where tissues close to veins remain green and rest of the lamina surface turns yellow.
6. **Leaf curl:** is curling of the leaves as a result of over growth on one side of the organ.
7. **Phyllody:** it is a metaplastic symptom where all the floral parts develop into leaf-like structures.
8. **Canker:** A necrotic, often sunken, lesion on a stem, branch, or twig of a plant.
9. **Anthracnose:** A disease that appear as black sunken leaf, stem or fruit lesions, caused by fungi that produced their asexual spores in an acervulus.
10. **Damping off:** Destruction of the seedlings near the soil line, resulting in seedlings falling over on the ground
11. **Mottle:** A symptom in which small but numerous areas of discolouration, commonly chlorotic, irregularly shaped and without sharply defined boundaries, stand out against a background of a different tint.  
**Yellows:** Because of the reduction in chlorophyll synthesis the presence of carotene and xanthophylls becomes evident even in young leaves leading to yellowing.


## PHYLUM CHYTRIDIOMYCOTA AND OOMYCOTA

1. **Genus *Synchytrium*:** All the species of the genus produce galls on different parts of the plants, particularly on the roots. The important species is *Synchytrium endobioticum* which causes was disease of potato. On the potato tubers, the warts are more typical and conspicuous, sometimes covering the whole tuber larger conspicuous, sometimes covering the whole tuber and larger than the tuber itself.

2. **Genus – *Pythium* (Damping off):** **Mycelium** (Aseptate, branched, cottony white); **Sporangiophores** (Different from vegetative hyphae, erect, simple and bearing sporangia singly); **Sporangia** (Spherical or globose, sometimes filamentous or toruloid); **Oospores** (Thick walled, spherical, usually smooth and three layered and plerotic); **Important species-** *P. aphanidermatum*, *P. ultimum*, *P. graminicolum* (damping off disease)




3. **Genus – *Phytophthora*:** **Mycelium** (Aseptate, coenocytic, branched); **Sporangiophores** (of indeterminate growth, zig-zag, sympodially branched, nodulate (with nodular swellings)); **Important species** (*P. infestans* (Late blight of potato); **Sporangia** (Single celled, lemon shaped and papillate); **Oospores** (Spherical in shape, smooth walled and aplerotic).




### Differences between *Pythium* and *Phytophthora*

S. N	Point of differences	<i>Pythium</i>	<i>Phytophthora</i>
1.	Haustoria	Absent	Rudimentary
2.	Sporangiophore	Of determinate growth	In determinate growth
3.	Sporangia	Spherical	Lemon shaped and papillate
4.	Vesicle	present	Normally absent
5.	Zoospore formation	In the vesicle	In the sporangium
6.	Oospore	Plerotic type	Aplerotic type
7.	Germination of Oospore	By germ tube	By germ sporangium

4. **Genus – *Peronospora*** (Downy mildew): **Mycelium** (Aseptate, coenocytic, branched, hyaline, endophytic and intercellular); **Conidia** (Single celled, spherical or oval in shape and borne singly); **Branching** (**Sterigmata**-Dichotomous at acute angles. Last (ultimate) branch is changed into the sterigmata); **Oospores** (Long and pointed and bearing conidia singly); **Conidiophores** (Spherical and reticulate in *Peronospora parasitica* (downy mildew of Crucifers).; Arise from the stomatal openings. They are slender, long, 2/3 portion unbranched and only 1/3 portion is branched; **Important species** (*Peronospora parasitica* (downy mildew of Crucifers), *P. tabacina* (downy mildew of tobacco). *P. pisi* (downy mildew of pea).




5. **Genus–*Sclerospora***: **Mycelium** (Aseptate, coenocytic, branched, hyaline, endophytic and intercellular); **Sporangiophores** (arise from the stomatal openings. They are short and broader towards apex); **Branching** – Dichotomous or even trichotomous. Last branch is changed into the sterigmata); **Sterigmata** (Short and swollen and bearing sporangia singly); **Sporangia** (Borne singly, single celled and sometimes papillate also); **Oospores** (Irregular in appearance because the sporangial wall shrinks and touches the oosporic wall at several places); **Important species** (*Sclerospora graminicola*, which causes green ear disease of Bajra).



### Differences between *Peronospora* and *Sclerospora*

S. N	Point of differences	<i>Peronospora</i>	<i>Sclerospora</i>
1.	Conidiophore / Sporangiphore	Long and slender	Short and broader at apex
2.	Branching	Dichotomous	Dichotomous or even trichotomous
3.	Sterigmata	Long and pointed	Short and swollen
4.	Conidia / sporangia	Conidia are formed	Sporangia are formed
5.	Oospore	Spherical/regular in appearance	Irregular in appearance

6. **Genus – *Albugo*** (White blister/rust): **Mycelium** (Aseptate, coenocytic, branched, hyaline, intercellular with knob shaped haustoria); **Sporangiophores** (Club shaped (clavate), simple, forming palisade layer below the epidermis, lateranl wall thickened and laterally free, bearing sporangia in basipetal chains); **Sporangia** (Single celled, globose and produced in chains in basipetal succession and attached with each other with a gelatinous pad known as “disjuncto”); **Oospores** (Rough and warty in appearance and yellow in colour); **Important species** (*Albugo candida* (white blister / white rust of crucifers).



### PHYLUM ZYGOMYCOTA

- Genus – *Mucor*** (Bread mould): **Mycelium** (Aseptate, branched, cottony white without stolons and rhizoids); **Sporangiophores** (Arise singly, simple, aseptate, bearing sporangia singly); **Sporangia** (Spherical or globose, smooth walled, fragile, columellate and multi-spored); **Columella** (Central portion in the sporangium which is sterile and “Dome shaped” ); **Aplanospores** (Oval or spherical in shape and single celled); **Zygosporos** (Rough walled, black, warty in appearance and provided with “suspensors”); **Important species** (*M. mucedo*, *M. basilliformis*).
- Genus – *Rhizopus*** (Bread mould): Characters of this genus are *Mucor*- like except the formation of stolons and rhizoids, sporangiophores arise in-groups from rhizoids); **Important species** (*R. stolonifera*).






### Differences between *Mucor* and *Rhizopus*

S. No.	Point of differences	<i>Mucor</i>	<i>Rhizopus</i>
1	Stolon	Absent	Present
2	Rhizoids	Absent	Present
3	Sprangiophores	Arise single	Arise in groups from rhizoids
4	Aplanospores	Simple	Striate (marked with lines)



### PHYLUM BASIDIOMYCOTA

- Genus – *Sphacelotheca*: Sorus** (Conical or cylindrical covered with the peridium and filled with black spore powder); **Columella** (In the central portion of sorus, slender on curved, made up of host tissues in *S. sorghi*); **Teliospores** (Round to shortly oval, dark brown in mass but olive brown singly, smooth walled. Mass but olive brown singly, smooth walled); Important spp. (*S. Sorghi* (Grain smut of Jowar), *S. cruenta* (Loose smut of jowar), *S. reiliana* (Head smut of Jowar)).
- Genus – *Tolyposporium*: Sorus** (Though formed in various parts of the host, is more common in the ovary); **Teliospores** (They are formed in the form of “spore balls” which are covered by member of host origin); Important species (*T. penicillariae* (smut of bajra), *T. ehrenbergii* (long smut of jowar))
- Genus- *Tilletia***: The disease caused by *Tilletia* are called as “Bunt”; **Teliospores** (Teliospores are large, 16-54 smooth, verrucose); Important species : *T. caries* & *T. foetida* (stinking smut or hill bunt)
- Genus – *Neovossia***: Grains partially or wholly converted into black powdery mass enclosed by membrane (*N. indica*); **Teliospores** (Dark brown, spherical to oval with reticulations on the episore, which appear as curved spines); Important species: *N. indica* (Karnal bunt of wheat), *N. horrida* (Bunt of rice).
- Genus – *Ustilago*: Sorus**: The teliosorus without a peridium; the black dusty teliospores are covered by a membrane of host origin; **Teliospores**: Small globose to oval or elliptical less than 20 μm in diameter in most of the species the outer wall (episore) is minutely echinulate but sometimes smooth also (*U. hordei*); Important species (*U. segetum tritici* (*U. tritici*); *U. nuda* – (Loose smut of barley); *U. maydis* (corn smut); *U. scitaminea* (whip smut of sugarcane)).

### Teliospores of Rust Fungi

<i>Uromyces</i>	<i>Puccinia</i>	<i>Melampsora</i>
<ul style="list-style-type: none"> <li>•Teliospores are stalked</li> <li>•They are single celled</li> <li>•Apex of teliospores is thickened</li> </ul> 	Teliospores are bicelled They are stalked 	Teliospores single celled, They are sessile and cylindrical in shape Form layer below the epidermis 

### Difference between Uredial and Telial stage

Uredial Stage		Telial Stage	
1. Epidermis ruptured Uredospore 2. Uredospores stalked 3. Uredospores finely echinulate 4. Capitulate paraphyses also present		1. Epidermis intact (unbroken) 2. Teliospores sessile 3. They are single celled, cylindrical in shape 4. Teliospores form layer below epidermis	

### Phylum Ascomycota

#### Class: Eurotiomycetes

- Genus – *Aspergillus* (Black mould): Mycelium** (Well developed, branched, septate, hyaline and submerged in the substratum); **Conidiophores** (Arise from the “foot cell,” aseptate, simple, terminating into vesicle); **Sterigmata** (Two rows of the sterigmata are formed on the vesicle. Primary sterigmata are flat. Secondary sterigmata are bottle shaped); **Conidia** (Borne on secondary sterigmata in long basipetal chains. They are globose, single celled, and echinulate); **Important species** (*A. niger*, *A. flavus*, *A. fumigatus*); **Perfect Stage** (*Eurotium*).
- Genus – *Penicillium* (Blue / green mould): Mycelium** (Well developed, branched, septate, hyaline and submerged in the substratum); **Conidiophores** (Septate and branched without forming vesicle. Foot cells absent); **Sterigmata** (Single row of sterigmata is formed. They are peg like); **Conidia** (Borne on sterigmata in long basipetal chains. They are, single celled, globose to ovoid, smooth walled and resemble as “glass beads”); **Important species** (*P. notatum*, *P. chrysogenum*); **Perfect Stage** (*Talaromyces*).

### Differences between *Aspergillus* and *Penicillium*

Point of differences	<i>Aspergillus</i>	<i>Penicillium</i>
1. Foot cell	Present	Absent
2. Conidiophores	Simple, aseptate	Septate and branched
3. Vesicle	Present	Absent
4. Sterigmata	Two rows	One row
5. Conidia	Echimulate (spiny)	Smooth
6. Perfect stage	<i>Eurotium</i>	<i>Talaromyces</i>

### Class –Sordariomycetes

- Genus – *Fusarium*: Mycelium** (Septate, branched, pinkish brown in colour); **Sporodochia** (Spherical, oval or ovate); **Conidiophores** (Short, aseptate or septate, usually simple may be branched also bearing conidia singly); **Conidia** (Microconidia – usually single celled or bicelled; Macroconidia – many celled (2-7), sickle shaped and knobbed at the base); Chlamydospores (formed in mycelium and macroconidia); **Important species** (*F. oxysporum* (wilt diseases), *F. udum* (wilt of pigeonpea); **Perfect Stage** (*Gibberella* and *Nectria*).
- Genus *Claviceps* (Ergot)** : The genus *Claviceps*, causes the important disease “Ergot” particularly of the cereals and millets. Common species is *C. purpurea* (Ergot of rye). **Mycelium** (Septate and branched, destroying ovary tissues and replacing it by cottony white mycelial mat forming conidiophores bearing conidia at their tips); **Conidia** (Minute, oval and single celled forming “Honey dew” stage (Nectar like secretion); **Sclerotia** (Black, hard and variable in shape and actually the ovaries being destroyed and replaced by sclerotia); **Perithecia** (Flask-like, ostiolate); **Asci** (Several in a perithecium, and are elongated, cylindrical in shape) **Ascospores** (Formed 8 in number in each ascus, which are long and thread like); Important sp. *C. purpurea* (Ergot of rye), *C. microcephala* (Ergot of bajra).
- Genus – *Pyricularia*: Conidiophores** (Straight, septate (with 2-4 septa), slender and thickened at the base); **Conidia** (Pyriiform (pear shaped) to obclavate base rounded tapering at the apex, 2- septate (three celled), slightly darkened. One to many conidia may found on a single conidiophore); **Important sp.** *P. oryzae* (blast of paddy); **Perfect stage** (*Magnaporthe oryzae*).
- Genus – *Colletotrichum*: Mycelium** (Septate, light brown, branched); **Acervuli** (Cushion shaped and provided with sterile, hair like black structure setae on acervuli); **Conidiophores** (Short, aseptate and unbranched); **Conidia** (Single celled, falcate, often with oil globule); **Important species** (*F. calcatum* (red rot of sugarcane), *C. truncatum* (Anthracnose of pulses); **Perfect stage** (*Glomerella*, *Physalospora*).

### Dothideomycetes

- Genus – *Helminthosporium*: Conidiophores** (Straight or zig-zag having knee joints (geniculate); **Conidia** (Conidia are produced singly at the apex and at knee-joints of the conidiophores. They are cylindrical, multi-septate, mostly with rounded ends); **Important sp.** (*H. gramineum* (stripe disease of barley) and *H. oryzae* (brown spot of paddy); **Perfect stage** (*Cochliobolus* and *Pyrenophora*).
- Genus – *Alternaria*: Conidiophores** (Septate, simple or sometimes branched); **Conidia** (Conidia borne usually in chains (acropetal). Sometimes solitary also. Conidia are provided with cross as well as longitudinal or oblique septa (muriform). Conidia are also provided with beak, which may vary from very short to very long according to species); **Important sp.** (*A. solani* (early blight of potato), *A. brassicae* (Alternaria blight of crucifers), *A. triticina* (Leaf blight of wheat); **Perfect stage** (*Pleospora*).
- Genus – *Phoma***: *Phoma* is similar to *Phyllosticta*; infect we call the same fungus as *Phyllosticta* when it occurs on leaves and *Phoma* when it occurs on the stem or other parts.
- Genus – *Phyllosticta*: Mycelium** (Well developed, branched and septate); **Pycnidia** (They are mostly flask shaped, dark, having natural opening known as “Ostiole”. Conidia are produced in pycnidia); **Conidiophores** (Short and simple); **Conidia** (Single celled, spherical or oval in shape, hyaline and come out in “Cirrhus” from the ostiole); **Important species** (*P. cajani* (leaf spot of pigeonpea); **Perfect Stage** (*Mycosphaerella*).
- Genus – *Cercospora*: Conidiophores** (Straight or zig-zag having knee joint (geniculate); **Conidia** (Conidia are produced singly at the apex and at knee joints of the conidiophores. They are acicular, multi-septate, tip acute and base broad); **Important sp.** (*C. personata* and *C. arachidicola*, which cause tikka disease of groundnut); **Perfect stage** (*Mycosphaerella*).

### Class: Letiomycetes

- Genus – *Erysiphe* (Powdery mildew): Mycelium** (Septate, branched, hyaline, ectophytic); **Asexual stage** (**Conidiophores**– Arise singly, short, septate, straight and simple); **Conidia** (Single celled, barrel shaped, hyaline and formed in basipetal chains); **Sexual stage** (**Cleistothecia**- Spherical in shape, black, hard, without any natural opening

(closed) and provided with appendages); **Appendages** (Many, hypha like (myceloid); **Asci** (Several in a cleistothecium, clavate, with 2, 4 or 8 ascospores); **Ascospores** (Usually spherical or oval, single celled, hyaline and formed in asci); **Important species** (*E. graminis tritici* (powdery mildew of wheat), *E. polygoni* (powdery mildew of pea), *E. cichoracearum* (powdery mildew of cucurbits).

- Genus - Sclerotinia: Mycelium** (Septate and branched mostly white in colour); **Conidiophores** (long septate and branched); **Conidia** (oval or lemon shaped, single celled and formed in chains); **Sclerotia** (Black, hard, variable in shape); **Apothecia** (Long and stalked cup or disc shaped); **Asci** (Clavate, slightly thickened at apex, with paraphyses); **Ascospores** (8 in number in each ascus, single celled round, or elliptical or elongated); Important sp. (*S. sclerotiorum* causing root rot and white rot disease).


### Class: Taphrinomycetes

- Genus Taphrina (Leaf curl fungus):** Mainly the spp. of this genus causes the disease symptoms as leaf curl, puckering, pockets and witches' broom. The most important species is *T. deformans*, the cause of "Peach leaf curl"; **Mycelium** (Composed of septate hyphae, consisting of typically binucleate cells. These hyphae may be intercellular on sub-cuticular or may grow within the walls for the epidermal cells); **Asci** (Naked, (without forming any fruiting body (ascocarp), forming the layer of naked (Hymenium on the epidermis of the host, and each ascus having 8 ascospores); **Ascospores** (Eight in number, mostly located at upper portion of asci, single celled, round or ovoid); Important sp.-*T. deformans* (Peach leaf curl) *T. pruni* (plum pocket).

## STAINING AND IDENTIFICATION OF PLANT PATHOGENIC BACTERIA

### A. Smear preparation:

- Take a grease free dry slide.
- Sterilize the inoculating loop on a flame of a Bunsen burner.
- Transfer a loopful of culture (or the specimen) by sterile loop and make a smear at the center. Smear should not be very thin or very thick.
- Allow the smear to dry in the air.
- Fix the dry smear by passing the slide 3-4 times through the flame quickly with the smear side facing up.



### B. Gram-staining procedure:

- Place the slides on the staining rods.
- Cover the smear with crystal violet stain and leave for 1 minute.
- Wash carefully under running tap water.
- Flood the smear with Gram's iodine solution and leave for 1 minute.
- Drain off the iodine Wash the slide for the again in a gentle stream of tap water.
- Flood the slide with the decolorizing agent then wait for 20-30 seconds. This can also be done by adding a drop by drop to the slide until the decolorizing agent running from the slides runs clear.
- Gently wash the slide under running tap water and drain completely.
- Counterstain with safranin for and wait for about 30 seconds to 1 minute.
- Wash slide in a gentle and indirect stream of tap water until no color appears in the effluent and then blot dry with absorbent paper and observe under the microscope.

**Gram Positive:** Dark purple (*Bacillus*, *Nocardia*, *Clostridium*, *Propionibacterium*, *Actinomyces*, *Enterococcus*, *Corynebacterium*, *Listria*, *Lactobacillus*, *Gardnerella*, *Mycoplasma*, *Staphylococcus*, *Streptomyces*, *Streptococcus* etc.)

**Gram Negative:** Pale to dark red (*Escherichia*, *Helicobcater*, *Hemophilus*, *Neisseria*, *Klebsiella*, *Enterobacter*, *Chlamydia*, *Vibrio*, *Pseudomonas*, *Salmonella*, *Shigella*).

## TRANSMISSION OF VIRUSES

### I. Sap transmission

- Young TLCV affected leaves of tomato plants are washed with tap water and dried with blotting sheet
- Leaves are weighed. Usually 10g leaf is preferred.
- For Preparation of standard extract potassium phosphate buffer volume equal to the weight of leaves (V/W, 1:1), is added into a mortar and leaves are ground with the pestle. (So, 10ml buffer is added).

4. After thorough grinding, the whole leaf pulp is passed through double layers of muslin cloth to get filtered standard extract of the leaves. This is best accomplished by pressing the juice (extract) through the muslin cloth used to hold the extract. The expressed juice (sap), which will contain the infectious principle, is used as inoculum.
5. Only young rapidly growing plants of urdbean showing distinct primary leaf stage (7-8 days after emergence) should be selected for inoculation.
6. Carborandum powder is slightly sprinkled on the leaves of test plants. This can be best done using a small sterilized cotton swab previously just touched with the carborandum powder separately taken in test tube or Petri dish.
7. The primary leaves of the test plants are inoculated by rubbing the sap over the leaf surface with quick, gentle strokes. Best result is observed if primary leaves are inoculated both sides i.e., upper surface as well as down surface.
8. Keep inoculated plants with proper label in the glasshouse for observations.
9. Keep regular watch and make observations every alternate day. ULCV will be seen transmitted systemically on the third trifoliate first after about 13-16 days after inoculation.

## II) Transmission by grafting

### 1) Wedge grafting

1. Upper of healthy tobacco plants are chopped off with a blade.
2. A longitudinal incision is given at the top of the stem. This serves as stock. A moist cotton swab is put to keep the stock moist and flow of the sap continuous.
3. Tobacco plant infected with leaf curl virus is used as scion.
4. Infected tobacco twig is given two deep incisions at the base to form the wedge, all the leaves except two or three at the top are removed.
5. Scion is fixed between the longitudinally incised arms of the stock.
6. The joint is wrapped with a moist-cotton swab and wrapped and tied with a polythene ribbon to retain moisture.
7. The graft is covered with a glass chimney whose top is covered with a square piece of polythene so that graft can get necessary moisture for survival. Inner wall of the chimney is wetted with water.
8. Everyday chimney should be removed for sometime so that graft plants get air.

**2) Approach grafting:** Diseased and healthy plants should be on two separate pots. A cut is made on diseased as well as on healthy plants at same length of the stem. Cuts on both plants should be brought together and tied with polythene and the whole set of plants covered with bell jar. Inner wall of the bell jar should be covered with moist blotting paper to retain moisture.

### 3) Bud grafting:

1. Buds are obtained from an infected twig of citrus plant bearing mature buds
2. Buds are cut with a knife
3. Wood portion is removed carefully
4. On the healthy nursery plant of citrus first a transverse and then a longitudinal incision are (T-shaped cut) made, which should be very superficial.
5. Bud is put inside the bark flap of the plants in such a way that only the bud remains emerged and bark portion of the bud remain covered with the flap.
6. Flap of the bark are carefully separated from the wood with the knife.
7. Grafted portion of the stem is tied with a polythene strip bearing the bud exposed.

**Other types of transmission of virus:** Transmission by insect vectors and by dodder.

## PHANEROGAMIC PLANT PARASITES

Some of the species of higher plants are known to live parasitically on other plants. These parasitic plants produce flowers and seeds also. They attack valuable crops and trees causing considerable damage. Some of these parasites attack roots of host while others parasitize stem. Some are devoid of chlorophyll (total parasite) and entirely depend on host for nutrition while others have chlorophyll (semi parasite) but no true roots and obtain water and mineral constituent of food from the host. The common parasitic flowering plants can be grouped as follows:

### I. Stem parasites


- a) Total parasite - *Cuscuta* sp. (Dodder)
- b) Semi-parasite - *Dendrophthoe* sp. (Loranthus)

### II. Root parasites


- a) Total parasite - *Orobanch* sp. (Broom rape)
- b) Semi-parasite - *Striga* sp. (Witch's weed)

**DODDER (*Cuscuta* sp.):** This is a non-chlorophyllous, leafless, twining parasitic seed plant, which attaches its yellow, orange or pink, thread-like stem to stem or other parts of (cultivated or wild plants). Leaves are represented by minute scales. It sends minute root like organs (haustoria) to the host cortex, which serve as an anchor as well as organs of food absorption. It bears tiny, white, pink or yellow flowers in cluster. Clover, berseem, flax and many oilseed crops are commonly attacked. The common important species is *Cuscuta gronovii*, which attacks garden, ornamental and hedge plants.

**LORANTHUS** (*Dendrophthiae* sp.)- Stem Semi-parasite: It is a common parasite of fruit trees. The parasite attacks aerial parts of host trees. It is devoid of a true root system of its own and hence, is dependent on host for water and mineral. Leaves are leathery and evergreen and possess chlorophyll. The stem is thick, erect or flattened at the nodes and appear to arise in cluster at the point of attack. Flowers are borne in clusters. They are long and tubular in shape and greenish-white or red in colour. The infected area of host becomes swollen and forms attachment disc e.g., *Dendrophthae falcate*, is an important species.



**BROOMRAPE** (*Orobanche* sp.): Root Total parasite: Affecting tobacco, brinjal, tomato, cabbage, cauliflower, turnip and many other Solanaceous and Cruciferous plants. The parasite consists of stout, fleshy stem, 15-20 cm tall. Stem is pale yellow or brownish-red in colour and covered by small, thin and brown scaly leaves. Flowers appear in axil of scales and are white and tubular. A large number of parasitic stems may be seen. *Orobanche ramosa* is an important species



**WITCH'S WEED** (*Striga* sp.): Witchweed is a well-known root semi-parasite of sugarcane, cereals, maize and millets in India. The parasite is a small plant, 15-30 cm tall with bright green, slightly hairy stem and leaves. Leaves are narrow, long and in opposite pairs. The flowers are small and usually brick red or scarlet, although some may be yellowish-red, yellowish or almost white. The seeds are borne in a capsule and are very minute to see with naked eye. Infected roots bear a large number of witch's weed haustoria, which are attached to root to feed on it. *Striga asiatica* is an important species.

## FUNGICIDES AND THEIR FORMULATIONS

**Fungicide:** The word is derived from the latin word *caedo* – to kill and the first term is fungus. Therefore, a fungicide is any agency that has the ability to kill a fungus e.g., heat, acid, UV-rays, light etc. However, in general the fungicide is defined as those chemicals capable of preventing infection of living plants by phytopathogenic fungi. Similarly this term could be applied in the case of Bacterial disease as Bactericides and in the case of Nematode infection it is Nematicides.

**Fumigants:** These are volatile chemicals applied into confined spaces or into a soil, which produces gas that destroys weed seeds and microorganisms and acts as a soil sterilant. The most common soil sterilants are methyl bromide, methane, allyl alcohol, carbon disulphide, chloropicrin and tetrachlorethane. They are packed in special pressure-resistant containers.

**Fumigation:** The application of a fumigant for disinfestation of an area.

**Fungicidal Dispenser:** An individual who has been certified to engage in the retail sale of fungicides for a licensed dealer.

**Fungicidal:** Killing fungal spores or mycelium. Applicable to physical agents such as heat, ultraviolet light, x-rays, gamma-radiation etc., as well as to chemicals that are lethal at low concentrations.

**Fungicide Dealer:** A person or firm holding a license to retail fungicides.

**Fungicide, Applicator:** An individual who provides services involving the use or application of fungicides.

**Fungicide, Eradicator:** (1) (curative fungicide) A fungicide used to control disease after infection has occurred. (2) A fungicide applied to a substratum in which the fungus is already present.

**Fungicide, Protective:** A fungicide used to protect an organism against infection by a fungus.

**Fungicide, Residue:** Fungicide remaining on or in a plant.

**Fungicide, Systemic:** A fungicide, which is absorbed through a plant surface and is translocated away from the site of application.

**Fungistatic:** Certain chemicals may temporarily inhibit fungus spore germination without being lethal. They are known as fungistatic. So, preventing the growth of a fungus without killing it.

**Fungistat:** A substance preventing the growth of a fungus without killing it.

**Fungistatis (mycostasis):** The prevention of fungal growth. The effect is reversible; if the inhibitor is removed or diluted, growth is resumed, cf. Fungicidal. In a broad sense the term can be applied to the non-germination of fungal spores due to the presence of auto-inhibitors or inhibitors from another organism or the substratum.

## Formulations of fungicides:

- 1. Wettable powder** is a very common formulation for most of the fungicides, which is used for spray mixtures. The modern wettable powders are water-dispersible, which have the quality to wet easily and disperse well in water. They are also called as Water-Dispersible Powders (WDP). The active ingredient is incorporated, usually at the rate of 30-80%, with a finely ground inert dust (filler) such as Kaolin, a wetting agent and a suspending agent.
- 2. Dust formulations** usually contain 1-10% active ingredient for direct application in dry forms. They are manufactured in such a way that they are light enough to be carried by a slight breeze for a considerable distance. The finely divided particle of active ingredient is carried on a carrier particle. The commonly used carriers (diluent) are attapulgite, kaolin, talc, pyrophyllite, diatomaceous earth, bentonite, calcium silicate, hydrated silica, calcium carbonate, magnesium carbonate, gypsum, lime etc.
- 3. Water dispersible Powders (WDP).** The active ingredient is incorporated, usually at the rate of 30-80%, with a finely ground inert dust (filler) such as Kaolin, a wetting agent and a suspending agent. The commonly used suspending agents are sodium lignin sulphonate (Sulphite dye), methyl celluloses, polyvinyl acetate and aluminium silicate. In addition, spreader-sticker is sometimes desirable, especially on plants with glossy or waxy leaves. Agitation is generally necessary to keep uniform suspension.
- 4. Granules (Pellets)** are the formulations of the fungicide with inert materials formed into particles about the size of coarse sugar. The granules normally contain 3-10% of the active ingredient. Due to their size, the granules do not drift but have limited application being confined to soil and seed treatments. Granules have the advantage they can be measured in dry form more easily and accurately than dusts or wettable powders. These are formulation in which a dry form of the active ingredient is mixed with a liquid. Such formulations usually contain a high percentage of active ingredient similar to wettable powders. They are mixed with water for final use and require agitation. These are mostly used as seed dressers in seed processing companies.
- 5. Solutions** are formulations in which active ingredient or a combination of active ingredients and a solvent is dissolved in water solutions. This has the advantage of requiring no agitation after formulation is added in water.
- 6. Suspension or slurries** are formulation in which a dry form of the active ingredient is mixed with a liquid. Such formulations usually contain a high percentage of active ingredient similar to wettable powders. They are mixed with water for final use and require agitation. These are mostly used as seed dressers in seed processing companies.

## Preparation of fungicidal solutions

- 1. Bordeaux mixture:** One kg of copper sulphate is powdered and dissolved in 50 litres of water. Similarly, 1 kg of lime is powdered and dissolved in another 50 litres of water. Then copper sulphate solution is slowly added to lime solution with constant stirring or alternatively, both the solutions may be poured simultaneously to a third contained and mixed well.  
**Merits:** Its natural tenacity to the plants. Its relative cheapness. Its utility in controlling wide variety of diseases. Somewhat non-toxic to human beings and cattle.  
**Demerits:** Its phytotoxic nature on certain plants like paddy, apples, peaches etc. It causes delay in ripening of fruits. The preparation is not very much practicable under field conditions. It's corroding action on metallic containers of spraying equipment. It is very much useful against a number of diseases like downy mildews, bacterial citrus canker etc.
- 2. Bordeaux paste:** Bordeaux Paste consists of same constituents as that of Bordeaux mixture, but it is in the form of a paste as the quantity of water used is too little. It is nothing but 10 per cent Bordeaux mixture and is prepared by mixing 1 kg of copper sulphate and 1 kg of lime in 10 litres of water. The method of mixing solution is similar to that of Bordeaux mixture. Wound dresser used to protect the wounded portions, cut ends of trees etc., against the infection by fungal pathogens.
- 3. Burgundy mixture:** It is prepared in the same way as Bordeaux mixture, except the lime is substituted by sodium carbonate. So, it is called as 'Soda Bordeaux'. It was developed Burgundy (France) in 1887 by Mason. The usual formula contains 1 kg of copper sulphate and 1 kg of sodium carbonate in 100 litres of water. It is a good substitute for Bordeaux mixture and used in copper-sensitive crops.
- 4. Cheshunt compound:** It is compound usually prepared by mixing 2 parts of copper sulphate and 11 parts of ammonium carbonate. This formula was suggested by Bewley in the year 1921. The two salts are well powdered, mixed thoroughly and stored in an air tight container for 24 hours before being used. The ripened mixture is used by dissolving it in water at the rate of 3 g/litre. Mixture is dissolved in a little hot water and volume is made up with cold water and used for spraying.
- 5. Chaubattia Paste:** Chaubattia paste is another wound dressing fungicide developed by Singh in 1942 at Government Fruit Research Station, Chaubattia in the Almora. It is usually prepared in glass containers or chinaware pot, by mixing 800g of copper carbonate and 800g of red lead in litre of raw linseed oil or lanolin. This paste is usually applied to pruned parts of apple, pear and peaches to control several diseases. The paste has the added advantage that it is not easily washed away by rain water.

## CALCULATION OF FUNGICIDAL SPRAYS CONCENTRATIONS

### Preparation of fungicidal solutions

**Bordeaux mixture:** Preparation of 1% Bordeaux Mixture  
1% means 5:5:50 or 5 gm. Copper sulphate + 5 gm. of slaked lime + 500 ml. of water

Since 250 ml. water contains 2.5 gm. of copper sulphate  
Therefore, 1 ml water will contain  $2.5 / 250$   
so that, 100 ml. water will contain  $2.5 / 250 \times 100 = 1\%$ .

Water 500 ml.  -----  -----  -----  -----	CuSO <sub>4</sub> 5 gm.  .....  .....  .....	Quick Lime 5 gm.  ○○○○○○ ○○○○○○ ○○○○○○
----------------------------------------------------------------------	-------------------------------------------------------------------	-------------------------------------------------------

### Exercise: Dry and wet seed treatment of seed at recommended doses:

**Requirements:** 1. Balance 2. Weight Box 3. Seed treating drum or Wooden container.

<b>Doses:</b>	<b>Agrosan G.N.</b>	<b>0.3%</b>
	<b>Thiram</b>	<b>0.2%</b>
	<b>Captan</b>	<b>0.2%</b>

**Method:** The required fungicide is weighed in the balance and put into a seed-treating drum. Seed of a particular crop is also placed into it and thoroughly mixed by agitation.

**Calculation:** For 8 Kg seed to be treated with Agrosan G.N. 0.3%

Since 100 gm. seed needs 0.3 gm. fungicide  
Therefore, 1 gm. seed needs  $0.3 / 100$   
and 1000 gm. or 1 Kg. seed will need  $0.3 / 100 \times 1000 = 3$  gm.  
1 Kg. seed needs 3 gm. fungicide  
therefore, 8 Kg. seeds will require  $8 \times 3 = 24$  g. fungicide.

### Exercise: To prepare fungicidal solution for spraying of 1 hectare of different crops.

**Requirements:** 1. Balance 2. Weight Box 3. container 4. Fungicide 5. Sprayer

<b>Doses:</b>	<b>Sulfex</b>	<b>0.3%</b>
	<b>Indofil M 45</b>	<b>0.2%</b>
	<b>Benlate</b>	<b>0.1%</b>
	<b>Water</b>	<b>1 litre</b>

**Method:** Different doses of fungicides were actually weighed and fungicidal solutions were prepared by adding water. These are then sprayed over the crop.

**Note:** For 1 hectare of spray, the water requirement is 1000 litre.

**Calculation:** For Indofil M 45

Since 100 cc water require 0.2 gm. fungicide  
therefore, 1 cc. water require  $0.2 / 100$   
and 1000 cc. or 1 litre water would require  $0.2 / 100 \times 1000 = 2$  gm.  
1000 cc or 1 litre require 2 gm. of fungicide  
therefore, 1000 litre will require  $1000 \times 2 = 2000$  gm. fungicides or 2 kg. fungicide.  
2 Kg. fungicide is required for 1 hectare field for 1 spraying.

**Precaution:** For suspensions constant agitation is necessary to avoid settling down of fungicides.

## METHODS OF APPLICATION OF FUNGICIDES AND THEIR SAFE USE

The fungicides are generally applied either as seed treatment, soil drenching, spraying and dusting. Out of these seed treatment and spraying are carried over throughout the whole world. At present dusting is mostly discarded and soil drenching is followed in a selective measure in nurseries and in the case of nematode control.

**A. Seed Treatment:** This is very much essential since a large number of pathogenic fungi and bacteria, are either carried outside or inside the seed i.e. they are either extremely seed borne or internally seed borne. When the seed germinates the

pathogen also become active and cause pre emergence, damping off or rotting. Sometimes post emergence diseases are also seen. The seed treatment also inhibits the Rhizosphere microflora also. The purpose of seed treatment is to destroy seed borne fungi and bacteria causing seed decay seedling blights and smuts etc. The treatment of seed of each and every crop should be done as a routine practice. Since it is a cheapest method of control throughout the whole world.

There are two methods of seed treatment:

1. **Physical method:** i.e. hot water or solar treatment for the control of loose smut of wheat
2. **Chemical control:** In this case either one or a combination of different fungicides is used to control. A number of diseases of a particular crop. This is because there is no ideal fungicide available at present which could control a number of pathogens of a particular plant.

A fungicide could be applied to seeds in either small lots using earthen pots or they may be applied to large seed lots by seed treating drums, commonly known as seed dressers. In other cases slurry treatment is also recommended. In the case of internal seed borne pathogens the seed or propagative parts are dipped in a fungicidal or bactericidal solutions for a specific time, i.e., from 10 min to 12 hours. I.e. antibiotics and mercurial fungicides.

In the case of Leguminous seeds the seed should not be treated the mercurial fungicides since they adversely affects modulation in the roots. In certain oilseed crops systemic fungicides are also banned since it causes harmful effect to human being and cattle.

**B. Soil Treatment:** This is classified into two :

1. **Physical method:** This involves use of heat in the form of steam or hot water or electricity.
2. **Chemical method:** In this case volatile and non-volatile chemicals are used. This is again classified into four groups:
  - i. **Soil drenching:** Fungicides are made up with water at the recommended concentration as in the case of spraying and applied to the soil surface wither before or after plant emergence. The required quantity of fungicide is applied with a sprinkler. It should be applied in such a way that the fungicide should reach a depth of at least 10-15 cm.
  - ii. **Broadcasting:** Sometimes non-volatile fungicides mixed with soil or fertilizer is scattered over the field uniformly with hand. Then it should be mixed thoroughly with a suitable implement. This method involves consumption of a large quantity of fungicides.
  - iii. **Furrow application:** In this method fungicides are applied either as dust or granules or powders into the furrow at the planting time. This method consumes much lesser quantity of fungicide then broadcasting, therefore it is economical.
  - iv. **Fumigation:** This is mainly done to control nematodes. The chemicals are applied under pressure by means of a soil fumigator. This produces gas which distributes itself throughout the soil. A polythene sheet is required to cover the fumigated portion of the soil for a limited period. This method is usually restricted to small areas only.
  - v. **Spraying:** This is the most common and widely adopted method. Spraying is done on leaves stems, and fruits. W. P. is mostly used and the most common carrier is water. The dispersion of spray martial is achieved by its passage under pressure through nozzles of sprayer. Spraying is of two types viz; high volume and low volume. These two terms are defined as the amount of liquid in which a crop protection material is applied to a crop. When spray involves large quantity of liquid/ unit area, they are turned as high volume. 600 lits and above per ha are considered to be of high volume category. In the case of low volume sprays it is possible to cover one has with about 100 lits or less amount of water. In high volume sprays the drop size in likely to be in between 0.5 to 3 mm where as in low volume sprays the droplet size is in between 15-40  $\mu$ . The low volume sprays are carried by means of air-blast equipment. Sometimes aircraft application is also used. Where the droplet size is 35 – 70 $\mu$ . In high volume sprays knapsack sprayer or foot sprayers are usually used.
  - vi. **Dusting:** Dust is applied to leaves fruits stems as an alternative to spraying. The dry powder is dusted by means of dusters for covering host surface. Generally dusting is possible. Only in calm weather also a better protection of plants is obtained, if the dust is applied, when the plant is slightly wet due to rain.